arXiv:1612.04972v3 [math.DG] 26 Dec 2016

Willmore orbits for isometric Lie actions*

Ming Xu and Jifu Li†

Abstract

In this work, we studied the Willmore submanifolds in a closed compact connected Riemannian manifold which are orbits for the isometric action of a compact connected Lie group. We call them homogeneous Willmore submanifolds or Willmore orbits. The criteria for these special Willmore submanifolds is much easier than the general theory which requires the Euler-Lagrange equation. Our main theorem claims, when the orbit type stratification for the group action satisfies certain conditions, then we can find a Willmore orbit in each stratified subset. Some classical examples of special importance, like Willmore torus, Veronese surface, etc., can be interpreted as Willmore orbits and easily verified with our method. Our theorems provide a large number of new examples for Willmore submanifolds, as well as estimates for their numbers which are sharp in some classical cases.

Mathematics Subject Classification (2010): 53C30, 53B25.

Key words: Willmore functional; Willmore submanifold; Willmore orbit; stratification; relative Willmore functional.

1 Introduction

A Willmore submanifold M^n is defined as an immersed closed submanifold in a closed compact connected Riemannian manifold (N^{n+p}, g_N), which represents a critical point for the Willmore functional $W_{N,g_N}(\cdot)$ [2][21][24]. It has been intensively studied in Riemannian geometry, and many examples have been found [7][10][11][12][19]. Studying Willmore functionals and Willmore submanifolds in the unit spheres are particularly important for many reasons. Firstly, they are invariance under conformal or Mobious translations. Secondly, they are closely related with the structure and classification theories for isoparametric hypersurfaces. And lastly, they are directly related with the famous Willmore conjecture which has been studied for decades by many geometers [13][16][20][23], and finally solved in 2014 [14]. Notice there is a general version for the Willmore conjecture [8][10][18], which is still widely open.

In this work, we study these subjects with Lie symmetry. We assume the metric admits the isometric actions of a compact connected Lie group, and study Willmore submanifolds which are orbits for this group action. We will simply call them homogeneous Willmore submanifolds or Willmore orbits.

The motivation for studying homogeneous Willmore submanifolds is two-folded. On one hand, since the Lie method has been introduced to the study of geometric functionals, it has been very successful on finding new examples of homogeneous minimal

*Supported by NSFC(no. 11671212)

†Corresponding author. E-mail: ljfanhan@126.com

submanifolds [6], or homogeneous Einstein metrics [1][5][22]. So we are inspired to try similar ideas or techniques for the Willmore functional, and expect finding new examples of Willmore submanifolds. On the other hand, we observe some classical examples of Willmore submanifolds with special importance, like the Willmore torus and its generalizations with higher codimensions[10][11], the Veronese surface in $S^4(1)$ [3][9][11], the focal submanifolds $\mathbb{C}\mathrm{P}^3$ and $\widetilde{\mathrm{G}}_2(\mathbb{R}^5)$ in $S^9(1)$ [19], and many others, permit large symmetries, i.e. they can be interpreted as Willmore orbits.

There is a simple criteria for Willmore orbits, which does not need to apply the general Euler-Lagrange equation [4][17]. First we have the orbit type stratification $N = \coprod_{\alpha\in\mathcal{A}} N_\alpha$ for the G-action on N according to the isotropy group types. Then a G-orbit $G\cdot x$ is a Willmore orbit iff it represents a critical point for $W_{N,g_N}(\cdot)|_{N_\alpha/G}$, where N_α is the stratified subset containing $G\cdot x$, and N_α/G is a finite dimensional smooth manifold contained in the defining space for the Willmore functional, i.e. the moduli space $\mathcal{M}_{M,N}$ of all immersion maps from $M = G\cdot x$ to N (see Theorem 4.2). Computing critical points in the finite dimensional N_α/G is much easier a task than in the infinitely dimensional $\mathcal{M}_{M,N}$.

The examples mentioned above can be easily verified to be Willmore submanifolds using this method. Further more, we expect finding new examples. The following theorem can meet out goal in quite general sense.

Theorem 1.1 *Let G be a compact connected Lie group acting isometrically on a closed compact connected Riemannian manifold (N, g_N). Denote $N = \coprod_{\alpha\in\mathcal{A}} N_\alpha$ the orbit type stratification for the G-action on N, where $\mathcal{A}$ is a finite set, and G/H_α the orbit type for each N_α. Assume the following two conditions:*

(1) $\dim H_\alpha < \dim G$ *for each* α*;*

(2) $\dim H_\alpha < \dim H_\beta$ *when* $N_\beta \subset \partial N_\alpha$.

Then there exists a Willmore orbit in each N_α.

The proof of Theorem 1.1 relies totally on our criteria Theorem 4.2 and the following theorem.

Theorem 1.2 *Let (N, g_N) and G be the same as in Theorem 1.1, and the orbit type stratification $N = \coprod_{\alpha\in\mathcal{A}} N_\alpha$ satisfy the same assumptions (1) and (2). Then for any sequence $x_n \in N_\alpha$, $n \in \mathbb{N}$, such that $\lim_{n\to\infty} x_n = x' \in N_\beta \subset \partial N_\alpha$, we have*

$$\lim_{n\to\infty} W_{N,g_N}([x_n]) = \infty,$$

in which $[x_n] \in N_\alpha/G \subset \mathcal{M}_{M,N}$, and M is any G-orbit in N_α.

Because Theorem 1.2 described the behavior of $W_{N,g_N}(\cdot)$ when the sequence of orbits $G\cdot x_n$'s collapse (converge) to one with a lower dimension, we will simply call it the orbit collapsing theorem for the Willmore functional.

Our methods are the standard localization technique in Riemannian geometry, i.e. enlarging a sequence of infinitesimal neighborhoods to a fixed standard model space

(see Section 6), together with the induction for the relative complexity degree between N_α and N_β (see Section 3 for its definition).

There are some tricky issues when we do the induction, because some extra product factor appears. Those extra product factors can not be easily "deleted" when the collapsing behavior of the orbits is complicated, for which we need to use the localization technique more than once. To avoid these subtle issues and make our argument rigorous, we switch to prove a more general theorem than Theorem 1.2, which includes the extra product factor. To be precise, we will define a relative Willmore functional, and prove a similar orbit collapsing theorem for our new functional (see Theorem 5.3). Then Theorem 1.2 and Theorem 1.1 are just immediate corollaries.

Imposing the assumptions (1) and (2) in Theorem 1.1 will not loss too much generalities. We can expect harvesting abundant new examples of Willmore submanifolds from Theorem 1.1. As examples, we will only discuss some important classical cases, i.e. the canonical action of $\mathrm{SO}(n_1) \times \mathrm{SO}(n_2)$ on $\mathbb{R}^{n_1+n_2+1}$ (and its generalization with more SO-factors), the $SO(3)$-conjugation on $S^4(1)$, and the Ad-action of $SO(5)$ on $S^9(1) \subset \mathfrak{g} = \mathfrak{so}(5)$. Notice Theorem 1.1 provides a lower bound estimate for the number of Willmore orbits, which is irrelevant to the equivariant changes for the metric g_N. For the examples we discuss, this estimate is sharp by either direct calculation or Theorem 5.1 in [10].

Notice some of above examples are discussed in [10] as isoparametric hypersurfaces. They were further studied in [19] to give more examples of non-homogeneous Willmore submanifolds, i.e. the focal submanifolds for an isoparametric foliation. We guess that some techniques in this work may be generalized to the isoparametric context and provide non-homogeneous Willmore submanifolds as well.

This paper is organized as following. In Section 2, we recall some fundamental knowledge about Willmore functional and Willmore submanifold. In Section 3, we introduce the orbit type stratification. In Section 4 we define homogeneous Willmore submanifold and prove its criteria, i.e. Theorem 4.2. In Section 5, we define the relative Willmore functional and state the orbit collapsing theorem for it, i.e. Theorem 5.3. In Section 6 and Section 7, we prove Theorem 5.3. In Section 8, we use our new method to reconsider some classical examples.

Acknowledgement. The authors would like to thank Shaoqiang Deng, Xiaobo Liu and Chao Qian for helpful discussions.

2 Willmore functional and Willmore submanifold

A closed compact connected n-dimensional immersed submanifold M in the $(n+p)$-dimensional closed compact Riemannian manifold (N, g_N) is called a *Willmore submanifold*, if it represents a critical point for the *Willmore functional*

$$W_{N,g_N}(M) = \int_M (S - n||H||^2)^{n/2} d\mathrm{vol}_{M,g_N}, \tag{2.1}$$

where S is the norm square of the second fundamental form of M in N, H is the mean curvature vector, and $d\mathrm{vol}_{M,g_N}$ is the volume form of M defined by the submanifold metric $g_M = i^* g_N$, where $i : M \to N$ is the immersion map [10].

The defining space of $W_{N,g_N}(\cdot)$ can be locally described as following. For each pair of closed compact connected manifolds (M^n, N^{n+p}), we define the moduli space of immersions from M to N as the quotient space of all immersion maps $i : M \to N$, divided by all self diffeomorphisms of M, and denote it as $\mathcal{M}_{M,N}$. A point in $\mathcal{M}_{M,N}$ can be either denoted as $[i]$ or $[i(M)]$, where i is the immersion map. Let g_N be a given Riemannian metric on N. Then for each immersion map i, the metric on M is naturally induced. So (2.1) gives the definition of the Willmore functional $W_{N,g_N}(\cdot)$ for each immersion. Obviously, it is preserved by all actions of self diffeomorphisms of M. So $W_{N,g_N}(\cdot)$ is well defined on $\mathcal{M}_{M,N}$.

Generally speaking, $\mathcal{M}_{M,N}$ is an infinitely dimensional manifold, if it is not empty. With the metric g_N chosen, the tangent space $T_{[i]}\mathcal{M}_{M,N}$ can be identified with the space $C^\infty(i^*\mathcal{V}^{i(M),N,g_N})$ of all smooth sections of the bundle $i^*\mathcal{V}^{i(M),N,g_N}$ over M, where $\mathcal{V}^{i(M),N,g_N}$ is the g_N-orthogonal complement sub-bundle for i_*TM in TN.

The tangent map (i.e. the first degree differential) DW_{N,g_N} of the Willmore functional is defined everywhere. An immersion map $i : M \to N$ defines a Willmore submanifold iff the linear functional $DW_{N,g_N} : T_{[i]}\mathcal{M}_{M,N} \to \mathbb{R}$ vanishes identically. This condition can be equivalently presented by the geometric data, which is called the Euler-Lagrange equation for the Willmore functional. See [4][11][17] for its explicit formula.

When $n = 0$ or 1, $W_{N,g_N}(\cdot)$ is a constant functional, and when $p = 0$, $\mathcal{M}_{M,N}$ is a discrete set. In this work, we count all immersed submanifolds in these special cases as Willmore submanifolds.

We have mentioned some classical examples of Willmore submanifolds with special importance in Section 1. We will revisit them in the last section, with more details given.

3 The orbit type stratification

Consider a closed compact connected Riemannian manifold N with the isometric action of a compact connected Lie group G. By the slice theorem in differential geometry [15], we have the *orbit type stratification* $N = \coprod_{\alpha\in\mathcal{A}} N_\alpha$, where $\mathcal{A}$ is a finite set. Here we summarize some fundamental properties for this stratification.

For each $\alpha \in \mathcal{A}$, N_α and N_α/G are connected smooth manifolds. They are not closed or compact in general. All G-orbits in a single N_α has the same isotropy group up to conjugations. If $G \cdot x = G/H_\alpha$ for some $x \in N_\alpha$, we will simply call H_α the *isotropy group type* of N_α, and call G/H_α the *orbit type* of N_α. The subset N_α is a smooth fiber bundle over N_α/G which fibers are G-orbits.

The closure $\overline{N_\alpha}$ of N_α in N is the union of N_α with several other N_β with lower dimensions. We denote $\partial N_\alpha = \overline{N_\alpha}\backslash N_\alpha$. For $N_\beta \subset \partial N_\alpha$, then we can find $x' \in N_\beta$ and $x \in N_\alpha$, such that the isotropy group H_β at x contains the isotropy group H_α at x'. Further more H_β either has a bigger dimension, or has the same dimension but more components than H_α.

Now we define the *relative complexity degree* between two stratified subsets N_α and $N_\beta \subset \partial N_\alpha$. It is the maximal integer d such that we can insert d different subsets N_{α_i}, $1 \le i \le d$, between $N_{\alpha_0} = N_\alpha$ and $N_{\alpha_{d+1}} = N_\beta$, such that $N_{\alpha_{i+1}} \subset \partial N_{\alpha_i}$ for $0 \le i \le d$.

In the rest of this section, we will construct a standard local chart at $x' \in N_\beta$, with respect to the group action and stratification.

For any $x' \in N_\beta$, the isotropy group H_β acts on $T_{x'}N$, which preserves the g_N-orthogonal decomposition

$$T_{x'}N = T_{x'}(G \cdot x') \oplus \mathcal{V}_{x'}^{N_\beta, N, g_N} \oplus \mathcal{V}_{x'}^{G\cdot x', N_\beta, g_N},$$

where the second and third factors in the right side are the g_N-orthogonal complement subspaces at x' for N_β in N and $G \cdot x'$ in N_β respectively. The H_β-action on the second factor will be important for our later discussions. We will simply call it the *isotropy action*.

We denote $n_1 = \dim G \cdot x'$, $n_2 = \dim N - \dim N_\beta$ and $n_3 = \dim N_\beta - \dim G \cdot x'$. The standard local chart is defined by the following diffeomorphism ϕ from $D^{n_1}(c_1) \times D^{n_2}(c_2) \times D^{n_3}(c_3)$ with some positive numbers c_1, c_2 and c_3, to a neighborhood $\mathcal{U}$ of x' in N, where $D^n(c) = \{v \in \mathbb{R} \text{ with } ||v|| < c\}$ denotes the open round disk centered at the origin o.

Firstly, we fix a linear direct sum decomposition $\mathfrak{g} = \mathfrak{h}_\beta + \mathfrak{m}$, where $\mathfrak{g} = \mathrm{Lie}(G)$ and $\mathfrak{h}_\beta = \mathrm{Lie}(H_\beta)$. Let $D^{n_1}(c_1)$ be an open disk around 0 in $\mathfrak{m}$ such that $\phi_1(v) = g(v) \cdot x$ with $v \in D^{n_1}(c_1)$ and $g(v) = \exp(v) \in G$, is a diffeomorphism from $D^{n_1}(c_1)$ to a neighborhood of x' in $G \cdot x$. Here exp is the exponential map in Lie theory.

Secondly, because the fixed point set $\mathrm{Fix}(H_\beta) \cap N_\beta$ is a smooth submanifold, we can choose a smooth normal slice $\mathcal{N}$ for $\mathrm{Fix}(H_\beta) \cap (G \cdot x')$ in $\mathrm{Fix}(H_\beta) \cap N_\beta$ such that $\mathcal{N}$ and $\mathrm{Fix}(H_\beta) \cap (G \cdot x')$ intersect g_N-orthogonally at x'. Schur Lemma for the H_β-actions implies $\mathcal{N}$ intersects g_N-orthogonally with $G \cdot x'$ at x', i.e. it is a g_N-orthogonal normal slice of $G \cdot x'$ in N_β. Let $D^{n_3}(c_3)$ be a local chart on this slice with x' identified with the origin o. Notice the isotropy group of each $v \in D^{n_3}(c_3) \subset \mathcal{N}$ is H_β.

Thirdly, we trivialize the bundle $\mathcal{V}^{N_\beta, N, g_N}$ around x' in $D^{n_3}(c_3) \in \mathcal{N}$ by n_2 smooth sections which provide a g_N-orthonormal basis at each point. Then any vector $v_2 \in D^{n_2}(c_2) \subset \mathcal{V}_{x'}^{N_\beta, N, g_N}$ can also be regarded as a tangent vector with the same g_N-length at any point in $D^{n_3}(c_3) \in \mathcal{N}$.

Finally we can construct the diffeomorphism as the following,

$$\phi(v_1, v_2, v_3) = g(v_1) \cdot \exp_{v_3, g_N} v_2, \tag{3.2}$$

where $v_i \in D^{n_i}(c_i)$ with sufficiently small positive c_i, $i = 1, 2, 3$. Here $\exp_{v_3, g_N}$ is the exponential map at $v_3 \in D^{n_3}(c_3) \subset N$ for the metric g_N.

We endow $D^{n_1}(c_1) \times D^{n_2}(c_2) \times D^{n_3}(n_3)$ with an Euclidean metric induced from g_N at x'. Obviously the three factors are then orthogonal to each other. We define the H_β-action on $D^{n_1}(c_1)$ and $D^{n_3}(c_3)$ to be trivial, and on $D^{n_2}(c_2)$ to be induced by the isotropy action. Then we have the following lemma.

Lemma 3.1 *When $x \in \mathcal{U}$ is sufficiently close to x', or c_1, c_2 and c_3 in above construction is sufficiently close to 0, we can find some $y \in D^{n_2}(c_2)$ and $z \in D^{n_3}(c_3)$, such that $D^{n_1}(c_1) \times (H_\beta \cdot y) \times z$ is an open neighborhood of x in $G \cdot x$. Further more, $H_\beta \cdot y$ is contained in $S^{n_2 - 1}(c)$ in the second factor $D^{n_2}(c_2)$ of $D^{n_1}(c_1) \times D^{n_2}(c_2) \times D^{n_3}(c_3)$, where c is the g_N-distance from x to N_β.*

The proof is not hard, by the construction for the diffeomorphism ϕ described above.

4 Willmore orbit and the criteria theorem

Let (N, g_N) be a closed compact connected Riemannian manifold, which admits the isometric action of a compact connected Lie group G. Denote $N = \coprod_{\alpha\in\mathcal{A}} N_\alpha$ the orbit type stratification, H_α and G/H_α the isotropy group type and orbit type for each N_α respectively.

Denote M any G-orbit in some N_α. Then N_α/G can be naturally identified as a submanifold of $\mathcal{M}_{M,N}$, with each $[x] \in N_\alpha/G$ for $x \in N_\alpha$ mapped to the element represented by the embedding $i : G \cdot x \to N$ in $\mathcal{M}_{M,N}$. Now we are ready to define homogeneous Willmore submanifolds.

Definition 4.1 *We will call the G-orbit $G \cdot x \subset N_\alpha$ a homogeneous Willmore submanifold or a Willmore orbit iff $[x] \in N_\alpha/G$ is the critical point for the Willmore functional $W_{N,g_N}(\cdot)$.*

The following theorem is a convenient criteria for Willmore orbits.

Theorem 4.2 *Let (N, g_N) be a closed compact connected Riemannian manifold with the isometric group action of a compact connected Lie group G. An orbit $G \cdot x$ in some N_α for the orbit type stratification of N is a Willmore orbit iff $[x] \in N_\alpha/G$ is a critical point for $W(\cdot)|_{N_\alpha/G}$.*

Proof. The proof for the "only if" part is trivial. We only need to prove the "if" part.

The isometric action of G on N can be naturally transported to $\mathcal{M}_{M,N}$. When M is a G-orbit in some N_α , the point $[x] \in N_\alpha/G \subset \mathcal{M}_{M,N}$ for any $x \in N_\alpha$ is a fixed point for this G-action. We want to decompose the tangent space $T_{[x]}\mathcal{M}_{M,N} = C^\infty(\mathcal{V}^{G\cdot x,N,g_N})$ into the sum of a sequence of finite dimensional irreducible representations. To make it precise, we first consider the Sobolev completion $\mathcal{H} = L^{2,k}(\mathcal{V}^{G\cdot x,N,g_N})$ of $T_{[x]}\mathcal{M}_{M,N}$. This is a Hilbert space with a countable basis. The action of G can be naturally extended to $\mathcal{H}$. If we define the $L^{2,k}$-norm as

$$||s||^2_{L^{2,k}} = \int_M (||s||^2 + ||\nabla s||^2 + \cdots + ||\nabla^k s||^2) d\mathrm{vol}_M,$$

using the norms $||\cdot||$ on sections induced by the G-invariant metric g_N, and G-invariant connections $\nabla : \Gamma(\mathcal{V}^{G\cdot x,N,g_N} \otimes \bigotimes^i T^*M) \to \Gamma(\mathcal{V}^{G\cdot x,N,g_N} \otimes \bigotimes^{i+1} T^*M)$, then the action of G preserves the inner product on $\mathcal{H}$. Because the G-action on N is isometric, it preserves the Willmore functional $W_{N,g_N}(\cdot)$ and its tangent map. When k is sufficiently big, by the Sobolev embedding theorem, the tangent map $DW_{N,g_N} : T_{[x]}\mathcal{M}_{M,N} \to \mathbb{R}$ of the Willmore functional can be extended to $\mathcal{H}$. So $DW_{N,g_N} : \mathcal{H} \to \mathbb{R}$ is a G-equivariant linear functional as well.

The trivial representations of G in $\mathcal{H}$ corresponds to all the G-invariant sections of $\mathcal{V}^{G\cdot x,N,g_N}$. All these sections give a finite dimensional subspace in $T_{[x]}\mathcal{M}_{M,N}$ which can be identified as the tangent space $T_{[x]}(N_i/G)$. So we can orthogonally decompose $\mathcal{H}$ as

$$\mathcal{H} = T_{[x]}(N_i/G) \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots ,$$

where each $\mathcal{H}_i$ is a finite dimensional nontrivial irreducible G-representation space.

By the assumption that $[x]$ is a critical point for $W_{N,g_N}(\cdot)|_{N_i/G}$, we have $DW_{N,g_N} = 0$ on $T_{[x]}N_i/G$. For each $\mathcal{H}_i$, $DW_{N,g_N}(\mathcal{H}_i) = 0$ by Schur Lemma. To summarize, DW_{N,g_N} at $[x]$ vanishes identically on $\mathcal{H}$. So $[x]$ is a critical point for the Willmore functional, i.e. $G \cdot x$ is a Willmore orbit. ■

An immediate consequence of Theorem 4.2 is the following corollary.

Corollary 4.3 *Let (N, g_N) be closed compact connected Riemannian manifold with the isometric group action of a compact connected Lie group G. Assume some subset N_α for the orbit type stratification is closed in N, then there exists a Willmore orbit in N_α. If N_α is not a single G-orbit, it contains at least two Willmore orbits.*

If N_α is a G-orbit, then it is obviously a Willmore orbit by Theorem 4.2. Otherwise we can find two Willmore orbits in N_α for the minimum and maximum points of $W_{N,g_N}(\cdot)|_{N_i/G}$.

5 The orbit collapsing theorem for the relative Willmore functional

We first define the relative Willmore functional.

Let N^{n+p} be an immersed or embedded submanifold of a Riemannian manifold $(N', g_{N'})$. Then we have the submanifold metric g_N on N induced by $g_{N'}$. Generally speaking we do not need particular assumptions for N', but we need N to be closed, compact and connected. Suppose M' is an immersed or embedded submanifold of N' which intersects with N transitively at a closed compact connected submanifold M^n.

Let $II_{M',g_{N'}}$ be the second fundamental form of M' in $(N', g_{N'})$, and $II_{M,M',g_{N'}}$ the restriction of $II_{M',g_{N'}}$ to M. Denote $S_{M,M',g_{N'}}$ and $H_{M,M',g_{N'}}$ the norm square and the trace vector of $II_{M,M',g_{N'}}$ with respect to the submanifold metric g_M induced by $g_{N'}$, and $d\mathrm{vol}_{M,g_{N'}}$ the volume form from g_M. Then the functional defined by

$$W_{N,N',g_{N'}}(M') = \int_M (S_{M,M',g_{N'}} - n||H_{M,M',g_{N'}}||^2)^{n/2} d\mathrm{vol}_{M,g_{N'}}$$

is called the *relative Willmore functional*.

We have the following obvious lemmas for the relative Willmore functional.

Lemma 5.1 *when $(N', g_{N'})$ is the Riemannian product of (N, g_N) with any manifold (D, g_D), N is identified with $N \times o$ for any fixed $o \in D$, and $M' = M \times D$ for a closed connected immersed submanifold M in N, then we have $W_{N,N',g_{N'}}(M') = W_{N,g_N}(M)$.*

Lemma 5.2 *For any constant $c > 0$, we have $W_{N,N',g_{N'}}(M') = W_{N,N',cg_{N'}}(M')$.*

The orbit collapsing theorem for relative Willmore functional is the following.

Theorem 5.3 *Let (N, g_N) and G be the same as in Theorem 1.1, and the orbit type stratification $N = \coprod_{\alpha \in \mathcal{A}} N_\alpha$ satisfy the same assumptions (1) and (2). Let (D, g_D) be any open set in $\mathbb{R}^q$ endowed with the standard Euclidean metric. Denote $g_{N',0}$ the product metric of g_N and g_D on $N' = N \times D$. Then for*

(1) *Any sequence of Riemannian metrics* $g_{N',n}$ *on* N'*, which converge to* $g_{N',0}$ *in the* C^∞*-topology,*

(2) *Any sequence* x_n *in the same stratified subset* N_α*, such that* $\lim_{n\to\infty} x_n = x' \in N_\beta \subset \partial N_\alpha$*,*

(3) *Any sequence* z_n *in* D *with* $\lim z_n = z \in D$*,*

we have

$$\lim_{n\to\infty} W_{N_n,N',g_{N',n}}(M'_n) = \infty,$$

where $N_n = N \times z_n$ *and* $M'_n = (G \cdot x_n) \times D$*.*

The proof of Theorem 5.3 will occupy the next two sections. Here we only use it to prove Theorem 1.2 and Theorem 1.1.

Proof of Theorem 1.2 Take a constant family of metrics $g_{N',n} = g_{N',0}$, and a constant sequence $z_n = z$, then Theorem 1.2 follows Lemma 5.1 and Theorem 5.3 immediately. ■

Proof of Theorem 1.1 Theorem 1.2 implies $W_{N,g_N}(\cdot)|_{N_\alpha/G}$ has a minimum point $[x]$ for some $x \in N_\alpha$. Then using Theorem 4.2, $G \cdot x$ is a Willmore orbit. ■

6 Proof of Theorem 5.3

The proof of Theorem 5.3 is an induction for the relative complexity degree between N_α and N_β.

Let (N, g_N) be a closed compact connected submanifold admitting the isometric actions of a compact connected Lie group G. Denote all the subsets for its orbit type stratification as N_α, $\alpha \in \mathcal{A}$, and assume they satisfy the conditions (1) and (2) in Theorem 5.3.

We will first prove Theorem 5.3 when the relative complexity degree between N_α and N_β is 0, which will occupy the rest of this section.

Denote $n_1 = \dim G \cdot x'$, $n_2 = \dim N - \dim N_\beta$, $n_3 = \dim N_\beta - \dim G \cdot x'$. We may assume n is sufficiently big, such that for each n, we can find a $x'_n \in N_\beta$, such that

$$d(x'_n, x_n) = d(x'_n, N_\beta) = r_n \to 0 \text{ when } n \to \infty,$$

where the distance function on N is induced by g_N. Then we have

$$x_n \in \exp_{x'_n,g_N} \mathcal{V}_{x'_n}^{N_\beta,N,g_N},$$

where the exponential map and the normal subspace at x is defined with respect to g_N. As we have shown in Section 3, we can construct a diffeomorphism ϕ as in (3.2) from an open neighborhood $\mathcal{U}$ of (x', z) in $N' = N \times D$, to the product $D^{n_1}(c_1) \times D^{n_2}(c_2) \times D^{n_3+q}(c_3)$, using the product metric $g_{N',0}$, where the open disks $D^{n_i}(c_i)$'s correspond to neighborhood of (x', z) in the G-orbit direction, the normal direction of N_β in N, and the normal direction of $(G \cdot x') \times z$ in $N_\beta \times D$ respectively.

When n is sufficiently large, the points x_n can be chosen from $\mathcal{U}$ such that $\phi^{-1}(x_n, z_n) = (y'_n, y''_n, y'''_n)$ with y'_n, y''_n and y'''_n all converging to the origin. Then we also have $\phi^{-1}(x'_n, z_n) = (y'_n, o, y'''_n)$. Define the metrics $g'_{N',n} = r_n^{-1} g_{N',n}$ on $N \times D$, and diffeomorphisms

$$\phi_n(v_1, v_2, v_3) = \phi(r_n v_1 + y'_n, r_n v_2, r_n v_3 + y'''_n).$$

Then for sufficiently large n, we can find a suitable neighborhood $\mathcal{U}_n$ of (x'_n, z_n) in $N \times D$, such that $\phi_n^{-1}(\mathcal{U}_n) = D^{n_1}(1) \times D^{n_2}(2) \times D^{n_3+q}(1)$, such that ϕ_n maps the origin to (x'_n, z_n), and the metrics $\tilde{g}_{\tilde{N}',n} = \phi_n^* g'_{N',n}$ on $\tilde{N}' = D^{n_1}(1) \times D^{n_2}(2) \times D^{n_3+q}(1)$ converges in the C^∞-topology, to the product metric $\tilde{g}_{N',0}$ of the standard Euclidean metrics on each factor.

Let $H_{\beta,o}$ be the identity component of H_β, and $y_n = r_n^{-1} y''_n$. Then by Lemma 3.1, $H_\beta \cdot y_n \subset S^{n_2-1}(1)$. Here we summarize some notations for the standard model space $\tilde{N}'$,

$$\begin{aligned}
\tilde{N}' &= D^{n_1}(1) \times D^{n_2}(2) \times D^{n_3+q}(1),\\
\tilde{S}' &= D^{n_1}(1) \times S^{n_2-1}(1) \times D^{n_3+q}(1),\\
\tilde{N}_n &= D^{n_1}(1) \times D^{n_2}(2) \times o,\\
\tilde{M}'_n &= D^{n_1}(1) \times (H_{\beta,o} \cdot y_n) \times D^{n_3+q}(1) \subset \tilde{S}',\\
\tilde{M}_n &= D^{n_1}(1) \times (H_{\beta,o} \cdot y_n) \times o = \tilde{N}_n \cap \tilde{M}'_n,\\
\tilde{g}_{\tilde{N}',n} &= \phi_n^* g'_{N',n}.\\
\tilde{g}_{\tilde{N}',0} &= \lim_{n\to\infty} \tilde{g}_{\tilde{N}',n}.
\end{aligned}$$

Notice $\tilde{M}'_n$, $\tilde{N}_n$ and $\tilde{M}_n$ are the pre-images of some neighborhoods of (x_n, z_n) in $M'_n = (G \cdot x_n) \times D$, $N_n = N \times z_n$ and $M_n = (G \cdot x_n) \times z_n$ respectively. By the assumption (2) in Theorem 5.3, $n_4 = \dim G \cdot x_n > n_1 = \dim G \cdot x'$, so $\dim \tilde{H}_{\beta,o} \cdot y_n = n_4 - n_1 > 0$. The metric $\tilde{g}_{\tilde{N}',0}$ is the product of standard Euclidean metrics on all factors.

The neighborhood $\mathcal{U}_n$ gets infinitely tiny when n goes to infinity. So for each n, we can find $l(n)$ different group elements $g_i \in G$, $1 \le i \le l(n)$, such that all $g_i \mathcal{U}_n$'s are disjoint with each other, and

$$\lim_{n\to\infty} \frac{l(n)}{n^{n_1}} > 0.$$

Now we prove for any n sufficiently big,

$$\int_{\tilde{M}_n} (S_{\tilde{M}_n, \tilde{M}'_n, \tilde{g}_{\tilde{N}',n}} - n_4 ||H_{M_n, M'_n, \tilde{g}_{\tilde{N}',n}}||^2)^{n_4/2} d\mathrm{vol}_{\tilde{M}_n, \tilde{g}_{\tilde{N}',n}} > C, \tag{6.3}$$

where C is a positive universal constant.

Let $\mathbf{n}_n$ and $\mathbf{n}$ be the outgoing unit normal vector fields of $\tilde{S}'$ in $\tilde{N}'$ with respect to $\tilde{g}_{\tilde{N}',n}$ and $\tilde{g}_{\tilde{N}',0}$ respectively. Because $\tilde{g}_{\tilde{N}',n}$ converges to $\tilde{g}_{\tilde{N}',0}$ in the C^∞-topology, $\mathbf{n}_n$ also converges to $\mathbf{n}$ in the C^∞-topology. The principal curvature of $\tilde{M}'_n$ in $\tilde{N}'$, for the normal vector field $\mathbf{n}$, with respect to $\tilde{g}_{\tilde{N}',0}$, equals 1 in the directions of the $H_{\beta,o} \cdot y_n \subset S^{n_2-1}(1)$, and equals 0 in the directions of $D^{n_1}(1)$. So when we change $\tilde{g}_{\tilde{N}',0}$ and $\mathbf{n}$ to $\tilde{g}_{\tilde{N}',n}$ and $\mathbf{n}_n$ respectively, those principal curvatures are still close to 1 and 0 respectively.

Assume $\lambda_{1,n} \leq \cdots \leq \lambda_{n_4,n}$ are all the eigenvalues of $II_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}(\mathbf{n}_n)$, i.e. the restriction of the second fundamental form of $\tilde{M}'_n$ to $\tilde{M}_n$, with respect to any local $\tilde{g}_{\tilde{N}',n}$-orthonormal tangent frame of $\tilde{M}_n$ and the normal vector field $\mathbf{n}_n$. Then when n is sufficiently big, we have $\lambda_{1,n} \leq 1/3$ and $\lambda_{n_4,n} \geq 2/3$. There exists a universal constant $C_1 > 0$, such that

$$\begin{aligned} & S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}} - n_4||H_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}||^2 \\ \geq & \sum_{i=1}^{n_4} \lambda_{i,n}^2 - \frac{1}{n_4}(\sum_{i=1}^{n_4} \lambda_{i,n})^2 > C_1. \end{aligned} \tag{6.4}$$

Notice all $H_{\beta,o} \cdot y_n$ belong to the same subset for the orbit type stratification of the $H_{\beta,o}$-action on $S^{n_2-1}(1)$. The orbits $H_{\beta,o} \cdot y_n$ in $S^{n_2-1}(1)$ can not have any collapsing subsequence, otherwise we can insert a stratified subset N_γ between N_α and N_β which is a contradiction that the relative complexity degree between N_α and N_β is 0. So there exists a positive lower bound for their volumes, with respect to the metric $\tilde{g}_{\tilde{N}',0}$. By the approximation of metrics, we also have

$$\mathrm{vol}(\tilde{M}_n, \tilde{g}_{\tilde{N}',n}) > C_2 > 0, \tag{6.5}$$

where C_2 is universal constant.

Summarize (6.4) and (6.5), we get

$$\int_{\tilde{M}_n} (S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}} - n_4||H_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}||^2)^{n_4/2} d\mathrm{vol}_{\tilde{M}_n,\tilde{g}_{\tilde{N}',n}} > C = C_1C_2 > 0,$$

which proves (6.3)

To finish the proof of theorem when the relative complexity degree between N_α and N_β is 0, we only need to use Lemma 5.2 and observe

$$W_{N_n,N',g_{N',n}}((G \cdot x_n) \times D) = W_{N_n,N',g'_{N',n}}((G \cdot x_n) \times D) > l(n) \cdot C \tag{6.6}$$

which diverges to infinity.

7 Proof of Theorem 5.3 continued

We continue to prove Theorem 5.3 with the assumption that the theorem is valid when the relative complexity degree between N_α and N_β is smaller than d.

Now we assume the relative complexity degree between N_α and N_β is d. We can apply exactly the same argument in the last section to reduce our discussion to the standard model $\tilde{N}'$, until we reach the step to prove (6.3), i.e.

$$\int_{\tilde{M}_n} (S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}} - n_4||H_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}||^2)^{n_4/2} d\mathrm{vol}_{\tilde{M}_n,\tilde{g}_{\tilde{N}',n}} > C > 0,$$

where C is a positive universal constant.

Passing to suitable subsequences, we may assume $y_n \subset S^{n_2-1}(1)$ converges to $y \in S^{n_2-1}(1)$. If $H_{\beta,o} \cdot y$ has the same dimension as $H_{\beta,o} \cdot y_n$, then (6.4) and (6.5) are

still valid, from which we can get (6.3). So we only need to consider the case that $\dim H_{\beta,o}\cdot y<\dim H_{\beta,o}\cdot y_n$.

We choose the local normal vector fields $\mathbf{n}_{i,n}$, $1\le i\le n_1+n_2-n_4$ on $\tilde{M}'_n$, with respect to $\tilde{g}_{\tilde{N}',n}$, satisfying the following conditions:

(1) $\mathbf{n}_{1,n}=\mathbf{n}_n$, and $\mathbf{n}_{i,n}$ is tangent to $\tilde{S}'$.

(2) For each n and at each point, all $\mathbf{n}_{i,n}$'s provide an $\tilde{g}_{\tilde{N}',n}$-orthonormal basis for the $\tilde{g}_{\tilde{N}',n}$-orthogonal complement bundle $\mathcal{V}^{\tilde{M}'_n,\tilde{N}',\tilde{g}_{\tilde{N}',n}}$.

The term

$$S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}-n_4||H_{M'_n,M_n,\tilde{g}_{\tilde{N}',n}}||^2$$

can be locally presented using $\mathbf{n}_{i,n}$. It is the sum of two parts. The first part $P_{1,n}$ contains all the terms for $\mathbf{n}_{1,n}=\mathbf{n}_n$, and $P_{2,n}$ contains all the terms for $\mathbf{n}_{i,n}$ with $i>1$. Both $P_{1,n}$ and $P_{2,n}$ are non-negative functions. In particular, because (6.4) is still valid when n is sufficiently big, i.e. $P_{1,n}>C_1>0$, then we have

$$\begin{aligned}&(S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}-n_4||H_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}||^2)^{n_4/2}\\ =\;&(P_{1,n}+P_{2,n})^{n_4/2}\ge C_2P_{1,n}^{n_1/2}P_{2,n}^{(n_4-n_1)/2}>C_1^{n_1/2}C_2P_{2,n}^{(n_4-n_1)/2},\end{aligned}\tag{7.7}$$

where C_2 is a positive universal constant.

Fix any $v_1\in D^{n_1}(1)$, we denote

$$\begin{aligned}\hat{N}'&=\tilde{S}'=D^{n_1}(1)\times S^{n_2-1}(1)\times D^{n_3+q}(1),\\ \hat{N}&=v_1\times S^{n_2-1}(1)\times o,\\ \hat{D}&=D^{n_1}(1)\times D^{n_3+q}(1),\\ \hat{M}'_n&=D^{n_1}(1)\times(H_{\beta,o}\cdot y_n)\times D^q(1),\\ \hat{M}_n&=v_1\times(H_{\beta,o}\cdot y_n)\times o,\\ \hat{g}_{\hat{N}',n}&=\tilde{g}_{\tilde{N}',n}|_{\hat{N}'},\\ \hat{g}_{\hat{N}',0}&=\lim_{n\to\infty}\hat{g}_{\hat{N}',n}\text{ in the }C^\infty\text{-topology.}\end{aligned}$$

Notice $\hat{g}_{\hat{N}',0}$ is just the Riemannian product of the standard metrics for all factors. The standard metric on the sphere factor is $H_{\beta,o}$-equivariant.

The complexity degree between the subset containing all $H_{\beta,o}\cdot y_n$ and that containing $H_{\beta,o}\cdot y$, for the orbit type stratification of the $H_{\beta,o}$-action on $S^{n_2-1}(1)$ is at most $d-1$, where d is the relative complexity between N_α and N_β. So by the inductive assumption, for each $v_1\in D^{n_1}(1)$, the term

$$\int_{\hat{M}_n}(S_{\hat{M}_n,\hat{M}'_n,\hat{g}_{\hat{N}',n}}-(n_4-n_1)||H_{\hat{M}_n,\hat{M}'_n,\hat{g}_{\hat{N}',n}}||^2)^{(n_4-n_1)/2}d\mathrm{vol}_{\hat{M}_n,\hat{g}_{\hat{N},n}}\tag{7.8}$$

goes to infinity with n.

Comparing the term

$$\hat{P}_n=S_{\hat{M}_n,\hat{M}'_n,\hat{g}_{\hat{N}',n}}-(n_4-n_1)||H_{\hat{M}_n,\hat{M}'_n,\hat{g}_{\hat{N}',n}}||^2$$

in (7.8) and the term $P_{2,n}$ in

$$S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}} - n_4||H_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}||^2,$$

containing all the terms for $\mathbf{n}_{i,n}$ with $i > 1$, we claim

Lemma 7.1 *For any $v_1 \in D^{n_1}(1)$ and at any point of $\hat{M}_n$, we have $\hat{P} \leq P_{n,2}$.*

Proof. Using the local orthogonal tangent frame $\{e_i, 1 \leq i \leq n_4\}$ on $\tilde{M}_n = D^{n_1} \times (H_{\beta,o} \cdot y_n) \times o$, with respect to $\tilde{g}_{\tilde{N}',n}$, such that all e_i with $i \leq n_4 - n_1$ are tangent to $\hat{M}_n = v_1 \times (H_{\beta,o} \cdot y_n) \times o$. At each point of $\hat{M}_n$, we can use this tangent frame to present $II_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}}(\mathbf{n}_{i,n})$ with $i > 1$ as an $n_4 \times n_4$ symmetric matrix $A_{i,n}$. Then its upper left $(n_4 - n_1) \times (n_4 - n_1)$-block $B_{i,n}$ is the matrix for $II_{\hat{M}_n,\hat{M}'_n,\hat{g}_{\hat{N}',n}}(\mathbf{n}_{i,n})$. So

$$P_{2,n} = \sum_{i>1} ||A_{i,n} - \frac{\mathrm{tr}A_{i,n}}{n_4} I||^2 = \sum_{i>1} \min_{c\in\mathbb{R}} ||A_{i,n} - cI||^2, \tag{7.9}$$

$$\hat{P}_n = \sum_{i>1} ||B_{i,n} - \frac{\mathrm{tr}B_{i,n}}{n_4 - n_1} I||^2 = \sum_{i>1} \min_{c\in\mathbb{R}} ||B_{i,n} - cI||^2, \tag{7.10}$$

where the norm square for a real matrix is the sum of the squares of all entries. Using the relation between $A_{i,n}$ and $B_{i,n}$ for each $i > 1$ and n to compare (7.9) and (7.10), we can easily get $\tilde{P}_n \leq P_{2,n}$ at each point of $\hat{M}_n$, and for any $v_1 \in D^{n_1}(1)$. ∎

Now we continue our estimate for (6.3). When n is sufficiently large,

$$\begin{aligned}
&\int_{\tilde{M}_n} (S_{\tilde{M}_n,\tilde{M}'_n,\tilde{g}_{\tilde{N}',n}} - n_4||H_{M'_n,M_n,\tilde{g}_{\tilde{N}',n}}||^2)^{n_4/2} d\mathrm{vol}_{\tilde{M}_n,\tilde{g}_{\tilde{N}',n}} \\
\geq\ & C_3 \int_{v_\in D^{n_1}(1)} \left[\int_{v_1\times\hat{M}_n} P_{2,n}^{(n_4-n_1)/2} d\mathrm{vol}_{\hat{M}_n,\hat{g}_{\hat{N}',n}} \right] \\
\geq\ & C_3 \int_{v_1\in D^{n_1}(1)} W_{\hat{N},\hat{N}',\hat{g}_{\hat{N}',n}}(\hat{M}'_n),
\end{aligned}$$

where C_3 is a positive universal constant.

By the inductive assumption, for each $v_1 \in D^{n_1}(1)$, the non-negative continuous function

$$f_n(v_1) = W_{\hat{N},\hat{N}',\hat{g}_{\hat{N}',n}}(\hat{M}'_n) = W_{\hat{N},\hat{N}',\hat{g}_{\hat{N}',n}}((H_{\beta,o} \cdot y_n) \times \hat{D})$$

goes to infinity with n. So

$$\lim_{n\to\infty} \int_{v_1\in D^{n_1}(1)} f_n(v_1) = \infty,$$

which is enough for us to get

$$\lim_{n\to\infty} W_{N_n,N',g_{N',n}}((G \cdot x_n) \times D) = \infty.$$

This ends the proof of Theorem 5.3, when the relative complexity between N_α and N_β is d. So by induction, the proof of Theorem 5.3 is done.

8 Some classical examples revisited

In this section, we revisit some important examples of Willmore submanifolds.

Example 1. The *Willmore torus* is the Willmore hypersurface $S^{n_1}(\sqrt{\frac{n_2}{n}})\times S^{n_2}(\sqrt{\frac{n_1}{n}})$ in S^{n+1}, where $n_1+n_2=n$ [10]. It is a principle orbit for the canonical action of $G=\mathrm{SO}(n_1+1)\times\mathrm{SO}(n_2+1)$ on $S^{n+1}(1)$. This group action is of cohomogeneity one, and the assumptions in Theorem 1.1 are satisfied. The explicit calculation (see Example 4 below) shows it is the only principal Willmore orbit predicted by Theorem 1.1.

Example 2. The *Veronese surface* is a $\mathbb{R}\mathrm{P}^2$ imbedded in $S^4(1)\subset\mathbb{R}^5$, which is a Willmore submanifold parametrized by $(x,y,z)\in\mathbb{R}^3$ with $x^2+y^2+z^2=3$ [3][9][11], i.e.

$$u_1=\frac{1}{\sqrt{3}}yz,\quad u_2=\frac{1}{\sqrt{3}}xz,\quad u_3=\frac{1}{\sqrt{3}}xy,$$
$$u_4=\frac{1}{2\sqrt{3}}(x^2-y^2),\quad u_5=\frac{1}{6}(x^2+y^2-2z^2).$$

Notice (x,y,z) and $(-x,-y,-z)$ define the same point $(u_1,\dots,u_5)$ on the Veronese surface. To see it is an orbit of a group action, we need to change the standard inner product to the following one. Identify $\mathbb{R}^5$ with the space of real trace-free symmetric 3×3-matrices, endowed with the inner product $\langle A,B\rangle=\mathrm{Tr}AB$. This inner product is preserved by all $\mathrm{SO}(3)$-conjugations. Denote $S^4(1)$ the unit sphere with respect to the new inner product. Then the conjugations of $G=\mathrm{SO}(3)$ on $S^4(1)$ is of cohomogeneity one. Its stratification consists of an open subset N_1 consisting of principal orbits, and two closed G-orbits N_2 and N_3. The two singular orbits are just the Veronese surfaces. By Corollary 4.3, we immediately see that they are Willmore orbits.

Theorem 1.1 predicts the existence of a principal Willmore orbit, which can be found the following observation. For any symmetric 3×3-matrix A, we denote its three eigenvalues as $\lambda_1(A)\le\lambda_2(A)\le\lambda_3(A)$. Then all the principal orbits M_t in N_1 can be parametrized by the value of $\lambda_2(\cdot)$, i.e. $t=\lambda_2(\cdot)\in(-1,1)$. The antipodal map on S^4 preserves $W(\cdot)$ and exchange M_t with M_{-t}. So $t=0$ is a critical point for $f(t)=W([M_t])$ with $t\in(-1,1)$. So by Theorem 4.2, M_0, the hypersurface of all symmetric matrices in $S^4(1)$ with vanishing determinants, is a Willmore orbit in $S^4(1)$.

Example 3. Consider the Ad-action of $G=\mathrm{SO}(5)$ on its Lie algebra $\mathfrak{g}$ preserves a canonical inner product $\langle\cdot,\cdot\rangle_{\mathrm{bi}}$. The unit sphere $S^9(1)$ of all vectors $v\in\mathfrak{g}$ with $\langle v,v\rangle_{\mathrm{bi}}=1$ is a cohomogeneity one space for this $\mathrm{SO}(5)$-action. The two singular orbits, i.e. $\widetilde{\mathrm{G}}_2(\mathbb{R}^5)=\mathrm{SO}(5)/\mathrm{SO}(2)\mathrm{SO}(3)$ and $\mathbb{C}\mathrm{P}^3=\mathrm{SO}(5)/\mathrm{U}(2)$, are Willmore submanifolds by Corollary 4.3. Theorem 1.1 predicts the existence of a principal Willmore orbit.

Example 4. Consider the the canonical action of $G=\mathrm{SO}(n_1+1)\times\cdots\times\mathrm{SO}(n_p+1)$ on $S^{n+p-1}(1)$ where $n_i>0$ for each i, and $\sum_{i=1}^p n_i=n$. The orbit type stratification for the G-action on N can be described as following. For each non-empty subset $A\subset\{1,\dots,p\}$, N_A is the union of all orbits

$$M_{t_1,\dots,t_p}=S^{n_1}(t_1)\times\cdots\times S^{n_p}(t_p),$$

where $\sum_{i=1}^p t_i^2=1$, and for each i, $t_i\ge 0$ and the equality happens iff $i\notin A$. Then $N=\coprod_{\emptyset\neq A\subset\{1,\dots,p\}}N_A$ is the orbit type stratification for the G-action on N. Obviously

$N_A \subset \bar{N}_B$ iff $A \subset B$. In particular, the open submanifold $N_{\{1,\dots,p\}}$ of N consists of all principal orbits $M_{t_1,\dots,t_p}$ with nonzero t_1, …, t_p.

Denote I_i and $\mathbf{n}_i$ the first fundamental form and the outgoing unit normal field of $S^{n_i}(1)$ in $\mathbb{R}^{n_i+1}$ respectively. Then restricted to $M_{t_1,\dots,t_p}$, the outgoing unit normal field of $S^{n+p-1}(1)$ in $\mathbb{R}^{n+p}$ can be presented as $\mathbf{n} = \sum_{i=1}^p t_i \mathbf{n}_i$.

The first fundamental form of $M_{t_1,\dots,t_p}$ is $I = t_1^2 I_1 + \cdots t_p^2 I_p$. Then second fundamental form $\widetilde{II}$ of $M_{t_1,\dots,t_p}$ in $\mathbb{R}^{n+p}$ is given by $\widetilde{II}(\mathbf{n}_i) = t_i I_i$. The mean curvature $\widetilde{H}$ of $\widetilde{II}$ satisfies $\widetilde{H}(\mathbf{n}_i) = \frac{n_i}{n} t_i^{-1}$. So with respect to the imbedding $M_{t_1,\dots,t_p} \subset S^{n+p-1}(1)$,

$$\begin{aligned} S &= \sum_{i=1}^p ||\widetilde{II}(\mathbf{n_i})||^2 - ||\widetilde{II}(\mathbf{n})||^2 = \sum_{i=1}^p n_i(t_i^{-2} - 1), \\ ||H||^2 &= ||\widetilde{H}||^2 - \widetilde{H}^2(\mathbf{n}) = \frac{1}{n^2}[\sum_{i=1}^p n_i^2 t_i^{-2}] - 1. \end{aligned}$$

So we have

$$\begin{aligned} f(t_1,\dots,t_p) &= W(M_{t_1,\dots,t_p}) \\ &= \int_{M_{t_1,\dots,t_p}} (S - n||H||^2)^{\frac{n}{2}} d\mathrm{vol}_{M_{t_1,\dots,t_p}} \\ &= C \cdot \prod_{i=1}^p t_i^{n_i} \cdot \left[\sum_{i=1}^p \frac{n_i(n-n_i)}{n} t_i^{-2}\right]^{\frac{n}{2}}, \end{aligned} \tag{8.11}$$

where C is a positive constant.

Notice $f(t_1,\dots,t_n)$ is homogeneous function of degree 0. So $M_{t_1,\dots,t_p} \subset S^n(1)$ is a Willmore orbit iff $(t_1,\dots,t_p)$ is a critical point for $f(t_1,\dots,t_p)$. The only critical point of f satisfying $\sum_{i=1}^p t_i^2 = 1$ and $t_i > 0$ for each i is given by $t_i = \sqrt{\frac{n-n_i}{n(p-1)}}$ for each i [11]. So the only principal Willmore orbit is

$$S^{n_1}(\sqrt{\frac{n-n_1}{n(p-1)}}) \times \cdots \times S^{n_p}(\sqrt{\frac{n-n_p}{n(p-1)}}). \tag{8.12}$$

Then we look for Willmore orbits in other N_A. The most obvious ones are N_A where A is a one-element set, which must be Willmore orbits by Corollary 4.3. In general the closure of any N_A is a unit sphere. So the above argument also proves there exists a unique Willmore orbit in N_A. For example, the only Willmore orbit in $N_{\{1,\dots,k\}}$ is

$$S^{n_1}(\sqrt{\frac{n'-n_1}{n'(k-1)}}) \times \cdots \times S^{n_k}(\sqrt{\frac{n'-n_k}{n'(k-1)}}) \times 0 \times \cdots \times 0,$$

where $n' = \sum_{i=1}^k n_i$.

To summarize, Theorem 1.1 provides sharp estimates for the number of Willmore orbits on the unit spheres in Example 1 and Example 4, which can be shown by short and explicit calculation. For the other two examples, we can apply Theorem 5.1 of [10] to show there are exactly three Willmore orbits, i.e. the estimates are still sharp, which can not be seen from Theorem 1.1. On the other hand, these estimates for the number of Willmore orbits are stable when the unit sphere metrics are equivalently perturbed.

References

[1] C. Böhm, Homogeneous Einstein metrics and simplicial complexes, J. Differential Geom. **67** (2004), 79-165.

[2] B. Y. Chen, Some conformal invariants of submanifolds and their applications, Boll. Un. Mat. Ital. **10** (1974), 380-385.

[3] S. S. Chern, M. do Carmo, S. Kobayashi, Minimal submanifolds of a sphere with second fundamental form of constant length, Functional Analysis and Related Fields, 59-75, Springer, New York, 1970.

[4] Z. Guo, H. Li and C. P. Wang, The second variantion formula for Willmore submanifolds in S^n, Results Math. **40** (2001), 205-225.

[5] J. Heber, Noncompact homogeneous Einstein spaces, Invent. Math. **133** (1998), 279-352.

[6] W. Y. Hsiang and H. B. Lawson, Minimal submanifolds of low cohomogeneity, J. Differential Geom. **5** (1971), 1-38.

[7] Z. Hu and H. Li, Willmore submanifolds in a Riemannian manifold, Proceedings of the workshop Contemporary Geometry and Related Topics, World Scientific, 251-275.

[8] O. Kobayashi, A Willmore type problem for $S^2 \times S^2$, Differential geometry and differential equations (Shanghai, 1985), 67-72, Lecture Notes in Math. **1255**, Springer, Berlin, 1987.

[9] H. B. Lawson, Local rigidity theorems for minimal hypersurfaces, Ann. of Math. (2) **89** (1989), 187-197.

[10] H. Li, Willmore hypersurfaces in a sphere, Asian J. Math. **5** (2001), 365-378.

[11] H. Li, Willmore submanifolds in a sphere, Math. Res. Lett., **9** (2002), 771-190.

[12] H. Li and G. Wei, Classification of Lagrangian Willmore submanifolds of the nearly Kaehler 6-sphere $S^6(1)$ with constant scalar curvature, Glasgow Math. J. **48** (2006), 53-64.

[13] P. Li and S. T. Yau, A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces, Invent. Math. **69** (1982), 269-291.

[14] F. C. Marques and A. Neves, Min-max theory and the Willmore conjecture, Ann. of Math. (2) **179** (2014), 683-782.

[15] D. Montgomery and C. T. Yang, The existence of a slice, Ann. of Math. (2) **65** (1957), 459-469.

[16] S. Montiel and A. Ros, Minimal immersions of surfaces by the first eigenfunctions and conformal area, Invent. Math. **86** (1986), 153-166.

[17] F. J. Pedit and T. J. Willmore, Conformal geometry, Atti Sem. Mat. Fis. Univ. Modena **36** (1988), 237-245.

[18] U. Pinkall, Inequalities of Willmore type for submanifolds, Math. Z. **193** (1986), 241-246.

[19] C. Qian, Z. Tang and W. Yan, New examples of Willmore submanifolds in the unit sphere via isoparametric functions, Ann. Global Anal. Geom. **42** (3) (2012), 403-410.

[20] A. Ros, The Willmore conjecture in the real projective space, Math. Res. Lett. **6** (1999), 487-493.

[21] C. P. Wang, Moebius geometry of submanifolds in S^n, Manuscripta Math. **96** (1998), 517-534.

[22] M. Y. Wang and W. Ziller, Existence and non-existence of homogeneous Einstein metrics, Invent. Math. **84** (1986), 177-194.

[23] T. J. Willmore, Notes on embedded surfaces, An. Sti. Univ. "A1. I. Cuza" Iasi Sect. I a Mat. (N. S.) **11B** (1965), 493-496.

[24] T. J. Willmore, Total Curvature in Riemannian Geometry, Ellis Horwood Limited, 1982.

Ming Xu
College of Mathematics, Tianjin Normal University, Tianjin 300387, P. R. CHINA,
Email: mgmgmgxu@163.com

Jifu Li
School of Science, Tianjin University of Technology, Tianjin 300384, P. R. CHINA,
Email:ljfanhan@126.com